

## **EXTENDED RANGE FORECAST OF ATLANTIC SEASONAL HURRICANE ACTIVITY AND LANDFALL STRIKE PROBABILITY FOR 2011**

We continue to foresee well above-average activity for the 2011 Atlantic hurricane season. We are predicting the same levels of activity that were forecast in early April due to the combination of expected neutral ENSO conditions and very favorable atmospheric and oceanic conditions in the tropical Atlantic. We continue to anticipate an above-average probability of United States and Caribbean major hurricane landfall.

(as of 1 June 2011)

By Philip J. Klotzbach<sup>1</sup> and William M. Gray<sup>2</sup>

This forecast as well as past forecasts and verifications are available via the World Wide Web at <http://hurricane.atmos.colostate.edu/Forecasts>

Emily Wilmsen, Colorado State University Media Representative, (970-491-6432) is available to answer various questions about this forecast

Department of Atmospheric Science  
Colorado State University  
Fort Collins, CO 80523  
Email: [amie@atmos.colostate.edu](mailto:amie@atmos.colostate.edu)

---

<sup>1</sup> Research Scientist

<sup>2</sup> Professor Emeritus of Atmospheric Science

ATLANTIC BASIN SEASONAL HURRICANE FORECAST FOR 2011

| Forecast Parameter and 1950-2000<br>Climatology (in parentheses) | Issue Date<br>8 December 2010 | Issue Date<br>6 April 2011 | Issue Date<br>1 June 2011 |
|--|-------------------------------|----------------------------|---------------------------|
| Named Storms (NS) (9.6)  | 17                            | 16                         | 16                        |
| Named Storm Days (NSD) (49.1)                                    | 85                            | 80                         | 80                        |
| Hurricanes (H) (5.9)   | 9                             | 9                          | 9                         |
| Hurricane Days (HD) (24.5)                                       | 40                            | 35                         | 35                        |
| Major Hurricanes (MH) (2.3)                                      | 5                             | 5                          | 5                         |
| Major Hurricane Days (MHD) (5.0)                                 | 10                            | 10                         | 10                        |
| Accumulated Cyclone Energy (ACE) (96.1)                          | 165                           | 160                        | 160                       |
| Net Tropical Cyclone Activity (NTC) (100%)                       | 180                           | 175                        | 175                       |

PROBABILITIES FOR AT LEAST ONE MAJOR (CATEGORY 3-4-5) HURRICANE  
LANDFALL ON EACH OF THE FOLLOWING COASTAL AREAS:

- 1) Entire U.S. coastline - 72% (average for last century is 52%)
- 2) U.S. East Coast Including Peninsula Florida - 48% (average for last century is 31%)
- 3) Gulf Coast from the Florida Panhandle westward to Brownsville - 47% (average for last century is 30%)

PROBABILITY FOR AT LEAST ONE MAJOR (CATEGORY 3-4-5) HURRICANE  
TRACKING INTO THE CARIBBEAN (10-20°N, 60-88°W)

- 1) 61% (average for last century is 42%)

2011 STATE IMPACT PROBABILITIES (NUMBERS IN PARENTHESES ARE  
LONG-PERIOD AVERAGES)

| State          | Hurricane | Major Hurricane |
|----------------|-----------|-----------------|
| Texas          | 50% (33%) | 20% (12%)       |
| Louisiana      | 47% (30%) | 20% (12%)       |
| Mississippi    | 18% (11%) | 8% (4%)         |
| Alabama        | 26% (16%) | 4% (3%)         |
| Florida        | 71% (51%) | 34% (21%)       |
| Georgia        | 19% (11%) | 2% (1%)         |
| South Carolina | 28% (17%) | 7% (4%)         |
| North Carolina | 44% (28%) | 13% (8%)        |
| Virginia       | 11% (6%)  | 1% (1%)         |
| Maryland       | 2% (1%)   | <1% (<1%)       |
| Delaware       | 2% (1%)   | <1% (<1%)       |
| New Jersey     | 2% (1%)   | <1% (<1%)       |
| New York       | 13% (8%)  | 6% (3%)         |
| Connecticut    | 12% (7%)  | 3% (2%)         |
| Rhode Island   | 10% (6%)  | 4% (3%)         |
| Massachusetts  | 12% (7%)  | 3% (2%)         |
| New Hampshire  | 2% (1%)   | <1% (<1%)       |
| Maine          | 7% (4%)   | <1% (<1%)       |

**Please also visit the Landfalling Probability Webpage at <http://www.e-transit.org/hurricane> for landfall probabilities for 11 U.S. coastal regions and 205 coastal and near-coastal counties from Brownsville, Texas to Eastport, Maine. In addition, we now include probabilities of named storms, hurricanes and major hurricanes tracking within 50 and 100 miles of various islands and landmasses in the Caribbean and Central America. We suggest that all coastal residents visit the Landfall Probability Webpage for their individual probabilities.**

## ABSTRACT

Information obtained through May 2011 indicates that the 2011 Atlantic hurricane season will have significantly more activity than the average 1950-2000 season. We estimate that 2011 will have about 9 hurricanes (average is 5.9), 16 named storms (average is 9.6), 80 named storm days (average is 49.1), 35 hurricane days (average is 24.5), 5 major (Category 3-4-5) hurricanes (average is 2.3) and 10 major hurricane days (average is 5.0). The probability of U.S. major hurricane landfall is estimated to be about 140 percent of the long-period average. We expect Atlantic basin Net Tropical Cyclone (NTC) activity in 2011 to be approximately 175 percent of the long-term average. We have maintained our early April forecast, due to the expected combination of neutral ENSO conditions and favorable atmospheric and oceanic conditions in the tropical Atlantic.

This forecast is based on a new extended-range early June statistical prediction scheme that utilizes 29 years of past data. Analog predictors are also utilized. Overall, conditions remain conducive for a very active hurricane season.

## Why issue extended-range forecasts for seasonal hurricane activity?

We are frequently asked this question. Our answer is that it is possible to say something about the probability of the coming year's hurricane activity which is superior to climatology. The Atlantic basin has the largest year-to-year variability of any of the global tropical cyclone basins. People are curious to know how active the upcoming season is likely to be, particularly if you can show hindcast skill improvement over climatology for many past years.

Everyone should realize that it is impossible to precisely predict this season's hurricane activity in early June. There is, however, much curiosity as to how global ocean and atmosphere features are presently arranged as regards to the probability of an active or inactive hurricane season for the coming year. Our new early June statistical forecast methodology shows strong evidence over 29 past years that improvement over climatology can be attained. **We would never issue a seasonal hurricane forecast unless we had a statistical model developed over a long hindcast period which showed significant skill over climatology.**

We issue these forecasts to satisfy the curiosity of the general public and to bring attention to the hurricane problem. There is a general interest in knowing what the odds are for an active or inactive season. One must remember that our forecasts are based on the premise that those global oceanic and atmospheric conditions which preceded comparatively active or inactive hurricane seasons in the past provide meaningful information about similar trends in future seasons. This is not always true for individual seasons. It is also important that the reader appreciate that these seasonal forecasts are based on statistical schemes which, owing to their intrinsically probabilistic nature, will fail in some years. Moreover, these forecasts do not specifically predict where within the Atlantic basin these storms will strike. The probability of landfall for any one location along the coast is very low and reflects the fact that, in any one season, most U.S. coastal areas will not feel the effects of a hurricane no matter how active the individual season is.

## **Notice of Author Changes**

**By William Gray**

The order of the authorship of these forecasts was reversed in 2006 from Gray and Klotzbach to Klotzbach and Gray. After 22 years (1984-2005) of making these forecasts, it was appropriate that I step back and have Phil Klotzbach assume the primary responsibility for our project's seasonal forecasts. Phil has been a member of my research project for the last eleven years and was second author on these forecasts from 2001-2005. I have greatly profited and enjoyed our close personal and working relationship. Phil has an office next to my office, and we talk nearly every day.

Phil is now devoting much more time to the improvement of these forecasts than I am. I am now giving more of my efforts to the global warming issue and in synthesizing my projects' many years of hurricane and typhoon studies.

Phil Klotzbach is an outstanding young scientist with a superb academic record. I have been amazed at how far he has come in his knowledge of hurricane prediction since joining my project in 2000. I foresee an outstanding future for him in the hurricane field. The success of the last three years of seasonal forecasts is an example. He is currently developing new seasonal and two-week forecast innovations that are improving our forecasts. Phil was awarded his Ph.D. degree in 2007. He is spending most of his time working towards better understanding and improving these Atlantic basin hurricane forecasts.

### Acknowledgment

This year's forecasts are funded by private and personal funds. We thank the GeoGraphics Laboratory at Bridgewater State College (MA) for their assistance in developing the United States Landfalling Hurricane Probability Webpage (available online at <http://www.e-transit.org/hurricane>).

The second author gratefully acknowledges the valuable input to his CSU seasonal forecast research project over many years by former graduate students and now colleagues Chris Landsea, John Knaff and Eric Blake. We also thank Professors Paul Mielke and Ken Berry of Colorado State University for much statistical analysis and advice over many years. We thank Bill Thorson for technical advice and assistance.

## DEFINITIONS AND ACRONYMS

Accumulated Cyclone Energy (ACE) - A measure of a named storm's potential for wind and storm surge destruction defined as the sum of the square of a named storm's maximum wind speed (in  $10^4$  knots<sup>2</sup>) for each 6-hour period of its existence. The 1950-2000 average value of this parameter is 96 for the Atlantic basin.

Atlantic Multi-Decadal Oscillation (AMO) – A mode of natural variability that occurs in the North Atlantic Ocean and evidencing itself in fluctuations in both sea surface temperature and sea level pressure fields. The AMO is likely related to fluctuations in the strength of the oceanic thermohaline circulation. Although several definitions of the AMO are currently used in the literature, we define the AMO based on North Atlantic sea surface temperatures from 50-60°N, 10-50°W.

Atlantic Basin – The area including the entire North Atlantic Ocean, the Caribbean Sea, and the Gulf of Mexico.

El Niño – A 12-18 month period during which anomalously warm sea surface temperatures occur in the eastern half of the equatorial Pacific. Moderate or strong El Niño events occur irregularly, about once every 3-7 years on average.

Hurricane (H) - A tropical cyclone with sustained low-level winds of 74 miles per hour (33 ms<sup>-1</sup> or 64 knots) or greater.

Hurricane Day (HD) - A measure of hurricane activity, one unit of which occurs as four 6-hour periods during which a tropical cyclone is observed or is estimated to have hurricane-force winds.

Madden Julian Oscillation (MJO) – A globally propagating mode of tropical atmospheric intra-seasonal variability. The wave tends to propagate eastward at approximately 5 ms<sup>-1</sup>, circling the globe in roughly 40-50 days.

Main Development Region (MDR) – An area in the tropical Atlantic where a majority of major hurricanes form, which we define as 10-20°N, 20-70°W.

Major Hurricane (MH) - A hurricane which reaches a sustained low-level wind of at least 111 mph (96 knots or 50 ms<sup>-1</sup>) at some point in its lifetime. This constitutes a category 3 or higher on the Saffir/Simpson scale.

Major Hurricane Day (MHD) - Four 6-hour periods during which a hurricane has an intensity of Saffir/Simpson category 3 or higher.

Multivariate ENSO Index (MEI) – An index defining ENSO that takes into account tropical Pacific sea surface temperatures, sea level pressures, zonal and meridional winds and cloudiness.

Named Storm (NS) - A hurricane, a tropical storm or a sub-tropical storm.

Named Storm Day (NSD) - As in HD but for four 6-hour periods during which a tropical or sub-tropical cyclone is observed (or is estimated) to have attained tropical storm-force winds.

Net Tropical Cyclone (NTC) Activity – Average seasonal percentage mean of NS, NSD, H, HD, MH, MHD. Gives overall indication of Atlantic basin seasonal hurricane activity. The 1950-2000 average value of this parameter is 100.

Saffir/Simpson Scale – A measurement scale ranging from 1 to 5 of hurricane wind and ocean surge intensity. One is a weak hurricane; whereas, five is the most intense hurricane.

Southern Oscillation Index (SOI) – A normalized measure of the surface pressure difference between Tahiti and Darwin. Low values typically indicate El Niño conditions.

Sea Surface Temperature – SST

Sea Surface Temperature Anomaly – SSTA

Thermohaline Circulation (THC) – A large-scale circulation in the Atlantic Ocean that is driven by fluctuations in salinity and temperature. When the THC is stronger than normal, the AMO tends to be in its warm (or positive) phase, and more Atlantic hurricanes typically form.

Tropical Cyclone (TC) - A large-scale circular flow occurring within the tropics and subtropics which has its strongest winds at low levels; including hurricanes, tropical storms and other weaker rotating vortices.

Tropical North Atlantic (TNA) index – A measure of sea surface temperatures in the area from 5.5-23.5°N, 15-57.5°W.

Tropical Storm (TS) - A tropical cyclone with maximum sustained winds between 39 mph (18 ms<sup>-1</sup> or 34 knots) and 73 mph (32 ms<sup>-1</sup> or 63 knots).

Vertical Wind Shear – The difference in horizontal wind between 200 mb (approximately 40000 feet or 12 km) and 850 mb (approximately 5000 feet or 1.6 km).

1 knot = 1.15 miles per hour = 0.515 meters per second

# **1 Introduction**

This is the 28th year in which the CSU Tropical Meteorology Project has made forecasts of the upcoming season's Atlantic basin hurricane activity. Our research team has shown that a sizable portion of the year-to-year variability of Atlantic tropical cyclone (TC) activity can be hindcast with skill exceeding climatology. This year's June forecast is based on a new statistical methodology derived from 29 years of past data. Qualitative adjustments are added to accommodate additional processes which may not be explicitly represented by our statistical analyses. These evolving forecast techniques are based on a variety of climate-related global and regional predictors previously shown to be related to the forthcoming seasonal Atlantic basin TC activity and landfall probability. We believe that seasonal forecasts must be based on methods that show significant hindcast skill in application to long periods of prior data. It is only through hindcast skill that one can demonstrate that seasonal forecast skill is possible. This is a valid methodology provided that the atmosphere continues to behave in the future as it has in the past.

The best predictors do not necessarily have the best individual correlations with hurricane activity. The best forecast parameters are those that explain the portion of the variance of seasonal hurricane activity that is not associated with the other forecast variables. It is possible for an important hurricane forecast parameter to show little direct relationship to a predictand by itself but to have an important influence when included with a set of 2-3 other predictors.

A direct correlation of a forecast parameter may not be the best measure of the importance of this predictor to the skill of a 3-4 parameter forecast model. This is the nature of the seasonal or climate forecast problem where one is dealing with a very complicated atmospheric-oceanic system that is highly non-linear. There is a maze of changing physical linkages between the many variables. These linkages can undergo unknown changes from weekly to decadal time scales. It is impossible to understand how all these processes interact with each other. No one can completely understand the full complexity of the atmosphere-ocean system. But, it is still possible to develop a reliable statistical forecast scheme which incorporates a number of the climate system's non-linear interactions. Any seasonal or climate forecast scheme should show significant hindcast skill before it is used in real-time forecasts.

## **2 June Forecast Methodology**

### **2.1 New June Statistical Forecast Scheme**

We have developed a new June statistical forecast model which we are using for the first time this year. This model has been built over the period from 1982-2010 to incorporate the most recent and reliable data that is available. It utilizes a total of four predictors. The new Climate Forecast System Reanalysis (CFSR) (Saha et al. 2010) has been completed from 1979-2009, while the NOAA Optimum Interpolation (OI) SST



(Reynolds et al. 2002) is available from 1982-present. The CFSR will begin to be updated in real-time later this year, but for the time being, we utilize the NCEP/NCAR Reanalysis for 2010 and 2011 values. This new model shows significant skill in predicting levels of Net Tropical Cyclone (NTC) activity over the 1982-2010 developmental period. The model correlates with NTC at 0.81 when all years are included in the model, while a drop-one cross-validation (jackknife) analysis yields a correlation with NTC of 0.72.

Table 1 displays cross-validated NTC hindcasts for 1982-2010 using the new statistical scheme, while Figure 1 displays observations versus cross-validated NTC hindcasts. We have correctly predicted above- or below-average seasons in 24 out of 29 hindcast years (83%). Our predictions have had a smaller error than climatology in 21 out of 29 years (72%). Our average hindcast error is 34 NTC units, compared with 55 NTC units for climatology. Figure 2 displays the locations of each of our predictors, while Table 2 displays the individual linear correlations between each predictor and NTC over the 1982-2010 hindcast period. All predictors correlate significantly at the 95% level using a two-tailed Student's t-test and assuming that each year represents an individual degree of freedom. The reader will note that we are incorporating a dynamical SST forecast from the European Centre for Medium-Range Weather Forecasts (ECMWF). Hindcast data provided by Frederic Vitart indicates that the ECMWF model has significant forecast skill for SSTs across the various Nino regions for September from a 1 May forecast date. We utilize the ECMWF ensemble mean prediction for September Nino 3 SSTs. Hindcast data from 1982-2010 show that the ECMWF forecast from 1 May correlates with observed September Nino 3 SSTs at 0.81. Table 3 displays the 2011 observed values for each of the four predictors in the new statistical forecast scheme.

Table 1: Observed versus early June cross-validated hindcast NTC for 1982-2010 using our new forecast scheme. Average errors for cross-validated hindcast NTC and climatological NTC predictions are given without respect to sign. Red bold-faced years in the “Hindcast NTC” column are years that we did not go the right way, while red bold-faced years in the “Hindcast improvement over Climatology” column are years that we did not beat climatology. The hindcast went the right way with regards to an above- or below-average season in 24 out of 29 years (83%), while hindcast improvement over climatology occurred in 21 out of 29 years (72%).

| Year           | Observed NTC | Hindcast NTC | Observed minus<br>Hindcast | Observed minus<br>Climatology | Hindcast improvement<br>over Climatology |
|----------------|--------------|--------------|----------------------------|-------------------------------|--|
| 1982           | 38           | 61           | -24                        | -62                           | 39                                       |
| 1983           | 31           | 51           | -20                        | -69                           | 49                                       |
| 1984           | 80           | <b>105</b>   | -25                        | -20                           | <b>-5</b>                                |
| 1985           | 106          | <b>87</b>    | 19                         | 6                             | <b>-13</b>                               |
| 1986           | 37           | 71           | -34                        | -63                           | 29                                       |
| 1987           | 46           | 63           | -18                        | -54                           | 37                                       |
| 1988           | 117          | 141          | -24                        | 17                            | <b>-7</b>                                |
| 1989           | 130          | 144          | -14                        | 30                            | 15                                       |
| 1990           | 100          | 146          | -46                        | 0                             | <b>-46</b>                               |
| 1991           | 58           | 47           | 11                         | -42                           | 32                                       |
| 1992           | 67           | 48           | 19                         | -33                           | 14                                       |
| 1993           | 52           | 50           | 2                          | -48                           | 46                                       |
| 1994           | 35           | 49           | -13                        | -65                           | 51                                       |
| 1995           | 222          | 228          | -6                         | 122                           | 115                                      |
| 1996           | 192          | 165          | 27                         | 92                            | 65                                       |
| 1997           | 54           | <b>145</b>   | -92                        | -46                           | <b>-45</b>                               |
| 1998           | 169          | 182          | -13                        | 69                            | 55                                       |
| 1999           | 182          | 155          | 27                         | 82                            | 55                                       |
| 2000           | 134          | 120          | 13                         | 34                            | 20                                       |
| 2001           | 135          | 152          | -16                        | 35                            | 19                                       |
| 2002           | 83           | 29           | 55                         | -17                           | <b>-38</b>                               |
| 2003           | 175          | 130          | 45                         | 75                            | 30                                       |
| 2004           | 232          | 125          | 107                        | 132                           | 25                                       |
| 2005           | 279          | 142          | 138                        | 179                           | 42                                       |
| 2006           | 85           | <b>163</b>   | -78                        | -15                           | <b>-63</b>                               |
| 2007           | 99           | <b>161</b>   | -62                        | -1                            | <b>-61</b>                               |
| 2008           | 162          | 176          | -14                        | 62                            | 49                                       |
| 2009           | 69           | 75           | -6                         | -31                           | 25                                       |
| 2010           | 196          | 226          | -30                        | 96                            | 66                                       |
| <b>Average</b> | <b>116</b>   | <b>116</b>   | <b> 34 </b>                | <b> 55 </b>                   | <b>+21</b>                               |

### Observed vs. June Model Jackknifed NTC




Figure 1: Observed versus early June jackknifed hindcast values of NTC for 1982-2010. The hindcast model explains 51% of the variance from climatology.

### New June Forecast Predictors




Figure 2: Location of predictors for our early June extended-range statistical prediction for the 2011 hurricane season.

Table 2: Linear correlation between each 1 June predictor and NTC over the 1982-2010 hindcast period.

| Predictor  | Correlation w/ NTC |
|--|--------------------|
| 1) April-May SST (15-55°N, 15-35°W) (+)                                  | 0.61               |
| 2) April-May 200 MB U (0-15°S, 150°E-120°W) (+)                          | 0.65               |
| 3) ECMWF 1 May SST Forecast for September Nino 3 (5°S-5°N, 90-150°W) (-) | -0.47              |
| 4) May SLP (20-40°N, 30-50°W) (-)  | -0.44              |

Table 3: Listing of 1 June 2011 predictors for the 2011 hurricane season. A plus (+) means that positive values of the parameter indicate increased hurricane activity.

| Predictor  | 2011 Forecast Value |
|--|---------------------|
| 1) April-May SST (15-55°N, 15-35°W) (+)                                  | +0.3 SD             |
| 2) April-May 200 MB U (0-15°S, 150°E-20°W) (+)                           | +1.6 SD             |
| 3) ECMWF 1 May SST Forecast for September Nino 3 (5°S-5°N, 90-150°W) (-) | +0.3 SD             |
| 4) May SLP (20-40°N, 30-50°W) (-)  | -0.9 SD             |

## 2.2 Physical Associations among Predictors Listed in Table 2

The locations and brief descriptions of the predictors for our early June statistical forecast are now discussed. It should be noted that all predictors correlate with physical features during August through October that are known to be favorable for elevated levels of hurricane activity. All of these factors are generally related to August-October vertical wind shear in the Atlantic Main Development Region (MDR) from 10-20°N, 20-70°W as shown in Figure 3.




Figure 3: Vertical wind profile typically associated with (a) inactive Atlantic basin hurricane seasons and (b) active Atlantic basin hurricane seasons. Note that (b) has reduced levels of vertical wind shear.

For each of these predictors, we display a four-panel figure showing linear correlations between values of each predictor and August-October values of sea surface temperature (SST), sea level pressure, 200 mb zonal wind, and 850 mb zonal wind, respectively. In general, higher values of SSTs, lower values of SLPA, anomalous westerlies at 850 mb and anomalous easterlies at 200 mb are associated with active Atlantic basin hurricane seasons. SST correlations are displayed using the NOAA Optimum Interpolation (OI) SST, while SLP, 850 mb, and 200 mb zonal wind correlations are displayed using the Climate Forecast System Reanalysis (CFSR).

Predictor 1. April-May SST in the Eastern Atlantic (+)

(15-55°N, 15-35°W)

Warmer-than-normal SSTs in the eastern Atlantic during the April-May period are associated with a weaker-than-normal subtropical high and reduced trade wind strength during the boreal spring (Knaff 1997). Positive SST anomalies in April-May are correlated with weaker trade winds and weaker upper tropospheric westerly winds, lower-than-normal sea level pressures and above-normal SSTs in the tropical Atlantic during the following August-October period (Figure 4). All three of these August-

October features are commonly associated with active Atlantic basin hurricane seasons, through reductions in vertical wind shear, increased vertical instability and increased mid-tropospheric moisture, respectively. Predictor 1 correlates quite strongly ( $\sim 0.6$ ) with NTC. Predictor 1 also strongly correlates ( $r = 0.65$ ) with August-October values of the Atlantic Meridional Mode (AMM) (Kossin and Vimont 2007) over the period from 1982-2010. The AMM has been shown to impact Atlantic hurricane activity through alterations in the position and intensity of the Atlantic Inter-Tropical Convergence Zone (ITCZ). Changes in the Atlantic ITCZ bring about changes in tropical Atlantic vertical and horizontal wind shear patterns and in tropical Atlantic SST patterns.

Predictor 2. April-May 200-mb zonal winds in the south-central tropical Pacific (+)

(0-15°S, 150°E-120°W)

Anomalous upper-level westerly zonal winds in the south-central tropical Pacific are typically associated with ongoing La Niña conditions and a strong Walker Circulation. The spring months are the climatologically favored time for ENSO events to transition from one phase to another (e.g., El Niño to La Niña or vice versa). If the atmosphere is strongly locked into the La Niña phase as evidenced by anomalously strong upper-level westerly winds, the odds of transitioning to an El Niño are reduced. Figure 5 shows that positive values of this predictor are also associated with favorable hurricane formation conditions in the tropical Atlantic, including above-average SSTs and below-average SLPs and zonal wind shear.

Predictor 3. ECMWF 1 May SST Forecast for September Nino 3 (-)

(5°S -5°N, 90-150°W)

The ECMWF seasonal forecast system 3 has shown skill at being able to forecast SST anomalies associated with ENSO several months into the future (Stockdale et al. 2011). ENSO has been documented in many studies to be one of the primary factors associated with interannual fluctuations in Atlantic basin and U.S. landfalling hurricane activity (Gray 1984, Goldenberg and Shapiro 1996, Bove et al. 1998, Klotzbach 2011), primarily through alterations in vertical wind shear patterns. The ensemble-averaged ENSO forecast for September values of the Nino 3 region from a 1 May issue date correlates with observations at 0.81. When the ECMWF model predicts cool SST anomalies for September, it strongly correlates with observed cool anomalies throughout the tropical Pacific associated with La Niña conditions, as well as reduced vertical wind shear, especially across the Caribbean (Figure 6).

Predictor 4. May SLP in the central Atlantic (-)


(20-40°N, 30-50°W)

Low pressure during the month of May in the central Atlantic is associated with reduced trade wind strength across the tropical Atlantic. This reduced trade wind strength


promotes reduced upwelling, mixing and enhances ocean current flow from the south, all of which feed back to promoting the development or sustenance of warm anomalies in the tropical Atlantic. These warm anomalies tend to persist throughout the peak months of the hurricane season (Figure 7). Also, upper-level easterly anomalies in the Caribbean are associated with low values of this predictor.

August-October Correlations w/ Predictor 1 (1982-2010)


(a) SST



(b) SLP



(c) 850 mb U



(d) 200 mb U





Figure 4: Linear correlations between April-May SST in the eastern Atlantic (Predictor 1) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 850 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). All four of these parameter deviations in the tropical Atlantic are known to be favorable for enhanced hurricane activity.




August-October Correlations w/ Predictor 2 (1982-2010)


(a) SST



(b) SLP



(c) 850 mb U



(d) 200 mb U






Figure 5: Linear correlations between April-May 200-mb zonal winds in the south-central tropical Pacific (Predictor 2) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 850 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). All of these parameter deviations over the tropical Atlantic and tropical Pacific tend to be associated with active hurricane seasons.

August-October Correlations w/ Predictor 3 (1982-2010)


(a) SST



(b) SLP



(c) 850 mb U



(d) 200 mb U






Figure 6: Linear correlations between a 1 May ECMWF SST forecast for September Nino 3 (Predictor 4) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 850 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). The predictor correlates very strongly with ENSO as well as vertical shear in the Caribbean. The correlation scale has been flipped to allow for easy comparison of correlations for all four predictors.

August-October Correlations w/ Predictor 4 (1982-2010)

(a) SST

(b) SLP



(c) 850 mb U

(d) 200 mb U





Figure 7: Linear correlations between May sea level pressure in the central Atlantic (Predictor 4) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 850 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). The correlation scale has been flipped to allow for easy comparison of correlations for all four predictors.

### 3 Forecast Uncertainty

One of the questions that we are asked regarding our seasonal hurricane predictions is the degree of uncertainty that is involved. Our predictions are our best estimate, but there is with all forecasts an uncertainty as to how well they will verify.

Table 4 provides our early June forecasts, with error bars based on one standard deviation of the 1982-2010 cross-validated hindcast error. We typically expect to see 2/3 of our forecasts verify within one standard deviation of observed values, with 95% of forecasts verifying within two standard deviations of observed values.

Table 4: Model hindcast error and our 2011 hurricane forecast. Uncertainty ranges are given in one standard deviation (SD) increments.

| Parameter                           | Hindcast Error (SD) | 2011 Forecast | Uncertainty Range – 1 SD<br>(67% of Forecasts Likely in this Range) |
|-------------------------------------|---------------------|---------------|---|
| Named Storms (NS)                   | 3.7                 | 16            | 12.3 – 19.7   |
| Named Storm Days (NSD)              | 21.1                | 80            | 58.9 – 101.1  |
| Hurricanes (H)                      | 2.1                 | 9             | 6.9 – 11.1  |
| Hurricane Days (HD)                 | 10.2                | 35            | 24.8 – 45.2   |
| Major Hurricanes (MH)               | 1.6                 | 5             | 3.4 – 6.6   |
| Major Hurricane Days (MHD)          | 5.3                 | 10            | 4.7 – 15.3  |
| Accumulated Cyclone Energy (ACE)    | 48                  | 160           | 112 – 212   |
| Net Tropical Cyclone (NTC) Activity | 48                  | 175           | 127 – 223   |

## 4 Caribbean-Specific Forecast Methodology

We have developed a forecast for the Caribbean that we debuted last year. Predictors for the Caribbean may be different than those used for the entire Atlantic basin. We intend to explore additional region-specific forecasts in the future, such as forecasts for the Gulf of Mexico or the higher latitude Atlantic.

We define the Caribbean to extend from 10-20°N, 60-88°W. This model attempts to predict seasonal levels of Accumulated Cyclone Energy (ACE) generated in the Caribbean. Through a combination of three predictors discussed in detail below, we can issue a forecast that shows significant levels of hindcast skill for the Caribbean basin. As is done with the overall basinwide forecast, this Caribbean forecast will also be updated in early August.

### 4.1 Caribbean Statistical Forecast Scheme

We have found that using three spring predictors, we can obtain early June hindcasts that show considerable skill over the sixty-year development period from 1949-2008.

This new scheme was created by evaluating three spring predictors using least-squared regression. By definition, least-squared regression tends to be conservative, and therefore, predicting large outlier events can be quite challenging. In order to help adjust for this challenge, the hindcasts from the linear regression model were adjusted to the final hindcast in the following manner:

The standardized value of each hindcast was calculated. These hindcasts were then adjusted to the final hindcast by multiplying by the standard deviation of the observations. Since the standard deviation of the observations is considerably larger than the standard deviation of the hindcasts, this aids in the forecast of outlying events. Any hindcasts that resulted in a negative ACE prediction were assigned a final ACE value of 0.

Our statistical scheme shows significant hindcast skill, explaining 43% of the variance ( $r = 0.66$ ) over the 1949-2008 period. Table 5 displays our early June hindcasts for 1949-2008 using the new statistical scheme, while Figure 8 displays observations versus ACE hindcasts. Our early June hindcasts have correctly predicted above- or below-average seasons in 47 out of 60 hindcast years (78%). These hindcasts have had a smaller error than climatology in 42 out of 60 years (70%). Our average hindcast error is 8.4 ACE units, compared with 12.5 ACE units for climatology. This scheme also shows considerable stability when broken in half, correlating at 0.60 from 1949-1978 and 0.71 from 1979-2008. This new scheme is also well-tuned to the multi-decadal active hurricane periods from 1950-1969 and 1995-2007 versus the inactive hurricane period from 1970-1994 (Table 6). As with our entire Atlantic forecast, the forecast skill for the Caribbean basin since 1980 has been particularly significant. Forecast improvement over climatology was notably high.

Figure 9 displays the locations of the early June Caribbean-specific predictors used in this scheme in map form. Table 7 lists the three predictors that are utilized for this year's June forecast.

Table 5: Observed versus hindcast Caribbean ACE for 1949-2008. Average errors for hindcast ACE and climatological ACE predictions are given without respect to sign. Red bold-faced years in the “Hindcast ACE” column (2) are years that we did not go the right way, while red bold-faced years in the “Hindcast improvement over Climatology” column (5) are years that we did not beat climatology. The hindcast went the right way with regards to an above- or below-average season in 47 out of 60 years (78%), while hindcast improvement over climatology occurred in 42 out of 60 years (70%).

| Year           | (1)<br>Observed ACE | (2)<br>Hindcast ACE | (3)<br>Observed minus<br>Hindcast | (4)<br>Observed minus<br>Climatology | (5)<br>Hindcast improvement over<br>Climatology |
|----------------|---------------------|---------------------|-----------------------------------|--------------------------------------|---|
| 1949           | 7                   | 8                   | 0                                 | -7                                   | 7   |
| 1950           | 16                  | 29                  | -13                               | 2                                    | <b>-11</b>                                      |
| 1951           | 24                  | 23                  | 1                                 | 9                                    | 8   |
| 1952           | 8                   | <b>23</b>           | -15                               | -7                                   | <b>-8</b>                                       |
| 1953           | 7                   | 4                   | 3                                 | -8                                   | 5   |
| 1954           | 40                  | 20                  | 20                                | 25                                   | 5   |
| 1955           | 48                  | 47                  | 1                                 | 34                                   | 33  |
| 1956           | 5                   | 11                  | -7                                | -10                                  | 3   |
| 1957           | 0                   | 2                   | -2                                | -15                                  | 12  |
| 1958           | 9                   | <b>21</b>           | -12                               | -5                                   | <b>-7</b>                                       |
| 1959           | 1                   | 13                  | -12                               | -14                                  | 1   |
| 1960           | 17                  | 26                  | -9                                | 2                                    | <b>-7</b>                                       |
| 1961           | 38                  | 35                  | 3                                 | 23                                   | 20  |
| 1962           | 0                   | <b>41</b>           | -41                               | -15                                  | <b>-26</b>                                      |
| 1963           | 28                  | 33                  | -4                                | 14                                   | 9   |
| 1964           | 21                  | 23                  | -2                                | 6                                    | 4   |
| 1965           | 0                   | 3                   | -3                                | -14                                  | 11  |
| 1966           | 22                  | 21                  | 2                                 | 8                                    | 6   |
| 1967           | 27                  | 15                  | 13                                | 13                                   | 0   |
| 1968           | 0                   | 8                   | -8                                | -14                                  | 6   |
| 1969           | 12                  | <b>18</b>           | -6                                | -3                                   | <b>-4</b>                                       |
| 1970           | 4                   | 12                  | -8                                | -11                                  | 3   |
| 1971           | 19                  | <b>0</b>            | 19                                | 5                                    | <b>-15</b>                                      |
| 1972           | 0                   | 0                   | 0                                 | -14                                  | 14  |
| 1973           | 0                   | 0                   | 0                                 | -14                                  | 14  |
| 1974           | 18                  | <b>3</b>            | 15                                | 4                                    | <b>-11</b>                                      |
| 1975           | 4                   | 0                   | 4                                 | -10                                  | 6   |
| 1976           | 1                   | 0                   | 1                                 | -14                                  | 13  |
| 1977           | 1                   | <b>21</b>           | -20                               | -14                                  | <b>-6</b>                                       |
| 1978           | 11                  | 14                  | -2                                | -3                                   | 1   |
| 1979           | 26                  | <b>12</b>           | 14                                | 11                                   | <b>-3</b>                                       |
| 1980           | 21                  | 17                  | 3                                 | 6                                    | 3   |
| 1981           | 4                   | <b>15</b>           | -12                               | -11                                  | <b>-1</b>                                       |
| 1982           | 0                   | 0                   | 0                                 | -15                                  | 15  |
| 1983           | 0                   | 0                   | 0                                 | -15                                  | 15  |
| 1984           | 3                   | 3                   | 0                                 | -11                                  | 11  |
| 1985           | 1                   | 0                   | 1                                 | -13                                  | 12  |
| 1986           | 1                   | 3                   | -2                                | -14                                  | 12  |
| 1987           | 6                   | 9                   | -3                                | -8                                   | 5   |
| 1988           | 42                  | 23                  | 19                                | 27                                   | 8   |
| 1989           | 16                  | 19                  | -2                                | 2                                    | 0   |
| 1990           | 8                   | <b>22</b>           | -14                               | -7                                   | <b>-7</b>                                       |
| 1991           | 0                   | 0                   | 0                                 | -15                                  | 14  |
| 1992           | 0                   | 0                   | 0                                 | -15                                  | 15  |
| 1993           | 4                   | 0                   | 4                                 | -11                                  | 7   |
| 1994           | 3                   | 0                   | 3                                 | -12                                  | 9   |
| 1995           | 23                  | 23                  | 0                                 | 9                                    | 8   |
| 1996           | 22                  | 17                  | 5                                 | 8                                    | 3   |
| 1997           | 3                   | 6                   | -4                                | -12                                  | 8   |
| 1998           | 44                  | 23                  | 21                                | 30                                   | 8   |
| 1999           | 24                  | 37                  | -12                               | 10                                   | <b>-3</b>                                       |
| 2000           | 13                  | <b>29</b>           | -16                               | -1                                   | <b>-15</b>                                      |
| 2001           | 23                  | 27                  | -4                                | 9                                    | 5   |
| 2002           | 6                   | 8                   | -2                                | -8                                   | 6   |
| 2003           | 6                   | <b>23</b>           | -17                               | -9                                   | <b>-9</b>                                       |
| 2004           | 51                  | 34                  | 17                                | 36                                   | 20  |
| 2005           | 58                  | 39                  | 19                                | 43                                   | 25  |
| 2006           | 4                   | <b>30</b>           | -25                               | -10                                  | <b>-15</b>                                      |
| 2007           | 47                  | 25                  | 22                                | 32                                   | 11  |
| 2008           | 22                  | 40                  | -18                               | 7                                    | <b>-11</b>                                      |
| 2009           | 4                   | <b>17</b>           | -13                               | -11                                  | <b>-2</b>                                       |
| 2010           | 22                  | 58                  | -36                               | 7                                    | <b>-29</b>                                      |
| <b>Average</b> | <b>15</b>           | <b>16</b>           | <b> 8.4 </b>                      | <b> 12.5 </b>                        | <b> 4.1 </b>                                    |

**Caribbean Basin ACE - Observations vs. 1 June Hindcast (1949-2008)**




Figure 8: Observed versus non-jackknifed hindcast values of Caribbean Basin ACE for 1949-2008. The statistical scheme correlates with observations at 0.66.

Table 6: Hindcast versus observed average ACE for active vs. inactive multi-decadal periods. Percentage differences from the climatological average (1949-2008) are in parentheses.

| <i>Years</i>            | <i>Average Hindcast ACE</i> | <i>Average Observed ACE</i> |
|-------------------------|-----------------------------|-----------------------------|
| 1949-1969<br>(Active)   | 20 (133%)                   | 16 (107%)                   |
| 1970-1994<br>(Inactive) | 7 (47%)                     | 8 (53%)                     |
| 1995-2008<br>(Active)   | 26 (173%)                   | 25 (167%)                   |

## June Caribbean Predictors




Figure 9: Location of spring predictors for our Caribbean statistical prediction for the 2011 hurricane season.

Table 7: Listing of 1 June 2011 Caribbean Basin predictors using the June statistical model for the 2011 hurricane season. A plus (+) means that positive values of the parameter indicate increased hurricane activity during the following year.

| Predictor                                      | 2011 Forecast Values |
|--|----------------------|
| 1) May SST (15-35°N, 15-30°W) (+)              | +0.6 SD              |
| 2) May 925 MB U (7.5°S-2.5°N, 160°E-160°W) (-) | -1.9 SD              |
| 3) April-May 200 MB V (5-25°N, 60-85°W) (+)    | +1.8 SD              |

The Caribbean model is calling for a very active hurricane season in 2011. Our forecast model is calling for an ACE of 46, which is nearly three times as much as the 1949-2008 average of 15 ACE units.

### 4.2 Physical Associations among Predictors Listed in Table 7

The locations and brief descriptions of the three spring predictors for our June Caribbean statistical forecast are now discussed. It should be noted that all three forecast parameters



correlate significantly with seasonal physical features that are known to be favorable for elevated levels of hurricane activity. Table 8 and 9 display correlations between each predictor and August-October-averaged sea surface temperature, sea level pressure, 200 mb zonal wind and 925 mb zonal wind in the Main Development Region (MDR) and in the Caribbean, respectively. Since many storms that generate large values of ACE in the Caribbean form in the MDR, one would expect that these predictors would correlate with physical features in both regions. Correlations that are significant at the 95% level using a two-tailed Student's t-test are highlighted in bold-faced type.

Table 8: Correlations between 1 June Caribbean predictors and August-October values of Main Development Region (10-20°N, 20-70°W) sea surface temperature, sea level pressure, 200 mb zonal wind and 925 mb zonal wind.

| Predictor                                      | SST         | SLP          | 200 MB U     | 925 MB U    |
|--|-------------|--------------|--------------|-------------|
| 1) May SST (15-35°N, 15-30°W) (+)              | <b>0.66</b> | <b>-0.40</b> | <b>-0.36</b> | <b>0.37</b> |
| 2) May 925 MB U (7.5°S-2.5°N, 160°E-160°W) (-) | -0.05       | <b>0.37</b>  | <b>0.40</b>  | -0.19       |
| 3) April-May 200 MB V (5-25°N, 60-85°W) (+)    | <b>0.30</b> | <b>-0.37</b> | <b>-0.40</b> | <b>0.47</b> |

Table 9: Correlations between 1 June Caribbean predictors and August-October values of Caribbean (10-20°N, 60-88°W) sea surface temperature, sea level pressure, 200 mb zonal wind and 925 mb zonal wind.

| Predictor                                      | SST         | SLP          | 200 MB U     | 925 MB U     |
|--|-------------|--------------|--------------|--------------|
| 1) May SST (15-35°N, 15-30°W) (+)              | <b>0.65</b> | <b>-0.51</b> | <b>-0.49</b> | <b>0.42</b>  |
| 2) May 925 MB U (7.5°S-2.5°N, 160°E-160°W) (-) | -0.03       | <b>0.30</b>  | <b>0.39</b>  | <b>-0.49</b> |
| 3) April-May 200 MB V (5-25°N, 60-85°W) (+)    | 0.25        | -0.25        | -0.23        | <b>0.34</b>  |

As was done with our Atlantic basin forecast, for each of these predictors, we display a four-panel figure showing linear correlations between values of each predictor and August-October values of sea surface temperature, sea level pressure, 200 mb zonal wind, and 925 mb zonal wind, respectively. In general, higher values of SSTA, lower values of SLPA, anomalous westerlies at 925 mb and anomalous easterlies at 200 mb are associated with active Caribbean basin seasons.

1. May SST in the Subtropical Atlantic (+)

(15-35°N, 15-30°W)

Above-normal sea surface temperatures (SSTs) in the eastern subtropical Atlantic are associated with a weakened Azores high and reduced trade wind strength during the late spring (Knaff 1997). This predictor is very strongly positively correlated ( $r = 0.65$ ) with Caribbean basin sea surface temperatures during August-October (Figure 10 and Table 9). Lower pressures are typically experienced throughout the Caribbean as well, implying a very favorable thermodynamic environment for tropical cyclone formation and intensification in this area during the climatologically most active portion of the hurricane season. This predictor also correlates favorably with reduced vertical wind

shear in the Caribbean as well as above-average SSTs, below-average SLPs and reduced vertical wind shear over the MDR.

Predictor 2. May 925 mb U in the Central Tropical Pacific (-)

(7.5°S-2.5°N, 160°E-160°W)

Enhanced trade wind strength in the central tropical Pacific is typically associated with a La Niña event. Stronger-than-normal trades encourage mixing and upwelling, promoting cooling. In addition, Kelvin wave propagation and associated warm water transport from the western Pacific to the eastern Pacific is inhibited by this stronger easterly flow. Consequently, strong correlations exist between values of this predictor and Nino 3.4 values. The correlation between this predictor and the May Nino 3.4 value is 0.60, the correlation between this predictor and the June-July Nino 3.4 values is 0.64, and the correlation between this predictor and the August-October Nino 3.4 value is 0.55. ENSO is a critical player in Caribbean tropical cyclone activity, as alterations in vertical shear caused by ENSO are more significant here than in any other part of the Atlantic basin (Klotzbach 2011). There is a significant negative correlation (-0.49) between this predictor and trade wind strength during August-October and a significant positive correlation (0.39) between this predictor and 200 mb wind strength during August-October over the Caribbean, indicating the considerable influence that this predictor has on the vertical shear values experienced over the Caribbean (Figure 11 and Table 9).

Predictor 3. April-May 200 MB V in the Southwest Atlantic (+)

(5-25°N, 60-85°W)

Anomalous southerly flow in the southwest Atlantic Ocean is associated with an anomalous upper-level trough and surface high pressure over South America, consequently driving strong trades over the tropical eastern Pacific and promoting La Niña conditions. Also, southerly flow at upper levels in the Caribbean is indicative of an upper-level anticyclone to the east. This upper-level anticyclone imparts anomalous easterly upper-level flow, weakening the climatological upper-level westerlies and thereby weakening vertical shear over both the tropical Atlantic as well as the Caribbean. It appears that this predictor's impacts are more strongly related to conditions over the MDR as opposed to the Caribbean, as its correlations are statistically significant with SST, SLP, 200 MB U and 925 MB U over the MDR (Figure 12 and Table 8), while displaying only a significant correlation with 925 MB U over the Caribbean (Figure 12 and Table 9).

## Predictor 1




Figure 10: Linear correlations between sub-tropical Atlantic SST (Predictor 1) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 925 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). All four of these parameter deviations are known to be favorable for enhanced hurricane activity.

## Predictor 2




Figure 11: Linear correlations between May 925 mb zonal wind in the central tropical Pacific (Predictor 2) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 925 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). All four of these parameter deviations are known to be favorable for enhanced hurricane activity. Zonal wind values have been multiplied by -1 to allow for easy comparison with Figure 1.

### Predictor 3




Figure 12: Linear correlations between April-May 200 mb meridional wind in the Caribbean (Predictor 3) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 925 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). All four of these parameter deviations are known to be favorable for enhanced hurricane activity.

## 5 Analog-Based Predictors for 2011 Hurricane Activity

Certain years in the historical record have global oceanic and atmospheric trends which are similar to 2011. These years also provide useful clues as to likely trends in activity that the forthcoming 2011 hurricane season may bring. For this early June extended range forecast, we determine which of the prior years in our database have distinct trends in key environmental conditions which are similar to current April-May 2011 conditions as well as expected August-October conditions. Table 10 lists our analog selections. We searched for years that were generally characterized by neutral or weak La Niña conditions and above-average tropical Atlantic and far North Atlantic SSTs during April-May. During the August-October period, we looked for years that were generally characterized by neutral ENSO conditions and above-average tropical Atlantic SST conditions.

There were five hurricane seasons since 1949 with characteristics most similar to those listed. These five years are 1951, 1981, 1989, 1996, and 2008. We anticipate that the 2011 hurricane season will have slightly more activity than what was experienced in the average of these five years, due to the very active season predicted by our new statistical model.

Table 10: Best analog years for 2011 with the associated hurricane activity listed for each year.

| Year                 | NS        | NSD       | H        | HD        | MH       | MHD       | ACE        | NTC        |
|----------------------|-----------|-----------|----------|-----------|----------|-----------|------------|------------|
| 1951                 | 10        | 57.75     | 8        | 36.25     | 5        | 8.25      | 137        | 148        |
| 1981                 | 12        | 66.00     | 7        | 22.50     | 3        | 3.75      | 100        | 113        |
| 1989                 | 11        | 66.00     | 7        | 31.75     | 2        | 9.75      | 135        | 130        |
| 1996                 | 12        | 78.50     | 8        | 41.00     | 5        | 14.25     | 177        | 182        |
| 2008                 | 16        | 88.25     | 8        | 30.50     | 5        | 7.50      | 146        | 162        |
| Average              | 12.2      | 71.3      | 7.6      | 32.4      | 4.0      | 8.7       | 139        | 147        |
| <b>2011 Forecast</b> | <b>16</b> | <b>80</b> | <b>9</b> | <b>35</b> | <b>5</b> | <b>10</b> | <b>160</b> | <b>175</b> |

## 6 ENSO

The moderate-to-strong La Niña conditions of this past winter have continued to weaken since early April. According to the Climate Prediction Center (CPC), SST anomalies in the central and eastern tropical Pacific have now warmed to the point where ENSO is in its neutral state. While the moderation of tropical Pacific SSTs have continued over the past couple of months, we do not expect to see a transition to El Niño conditions during the next several months. Upper-ocean heat content anomalies in the eastern and central Pacific (0-300 meters), which became positive towards the end of February have leveled off over the past 6-8 weeks. This indicates that there has been little translation of the very warm waters of the western Pacific warm pool eastward since early April (Figure 13).




Figure 13: Central and eastern tropical Pacific upper ocean (0-300 meters) heat content anomalies over the past year. Note the significant warming of these anomalies from December 2010 through March 2011 and the leveling off since then.

Currently, SSTs anomalies are slightly below average for most of the tropical central Pacific, with above-average SST anomalies recently observed only in the Nino 1+2 of the tropical eastern Pacific. Table 11 displays March and May SST anomalies for several Nino regions. All of the Nino regions have experienced considerable warming over the past two months, but we do not believe this warming trend will continue.

Table 11: March and May SST anomalies for Nino 1+2, Nino 3, Nino 3.4, and Nino 4, respectively. May-March SST anomaly differences are also provided.

| Region   | March SST Anomaly (°C) | May SST Anomaly (°C) | May – March SST Anomaly (°C) |
|----------|------------------------|----------------------|------------------------------|
| Nino 1+2 | -0.4                   | 0.6                  | +1.0                         |
| Nino 3   | -0.8                   | -0.1                 | +0.7                         |
| Nino 3.4 | -0.8                   | -0.3                 | +0.5                         |
| Nino 4   | -1.0                   | -0.3                 | +0.7                         |

There is considerable uncertainty as to what is going to happen with the current neutral ENSO event. The spring months are known for their ENSO predictability barrier. While we are nearing the end of this predictability barrier, considerable changes with ENSO often take place through June and July. Both statistical and dynamical models show improved skill by the end of May for the August-October period when compared with their skill at the end of March. However, there remains a very wide spread in both the statistical and dynamical model guidance for the August-October period (Figure 14). We find that, in general, the European Centre for Medium-Range Weather Forecasts (ECMWF) shows the best prediction skill of the various ENSO models. The correlation skill between a 1 May forecast from the ECMWF model and the observed September

Nino 3.4 anomaly is 0.82, based on hindcasts/forecasts from 1982-2010, explaining approximately 65% of the variance in Nino 3.4 SST. For reference, the correlation skill of a 1 March forecast from the ECMWF model was 0.71, indicating that approximately 15% additional variance can be explained by shortening the lead time of the forecast from 1 March to 1 May. The average of the various ECMWF ensemble members is calling for a September Nino 3.4 SST anomaly of approximately  $+0.2^{\circ}\text{C}$ , while only a few members call for weak El Niño conditions. Even if a weak El Niño were to develop, the currently-observed very favorable tropical Atlantic conditions would likely still lead to an active hurricane season, similar to what was observed in 1951, 1953, 1969, and 2004. About 80% of ensemble members from the 1 May ECMWF prediction call for either neutral or weak La Niña conditions through September (Figure 15).




Figure 14: ENSO forecasts from various statistical and dynamical models. Figure courtesy of the International Research Institute (IRI). There is a very wide spread in the model guidance for the August-October period, with several models calling for either El Niño or La Niña conditions and the rest calling for neutral conditions.






Figure 15: ECMWF ensemble model forecast for the Nino 3.4 region. Green circles represent the approximate centroid of the ensembles for each individual month.

Another reason why we believe that we are not likely to see a rapid transition to El Niño is that the atmosphere is generally responding like La Niña conditions are still in progress. Anomalously strong trade winds persist near the International Date Line (Figure 16). Stronger-than-normal trades help prevent eastward-propagating Kelvin waves that transport warm waters from the western tropical Pacific eastward. Despite a brief incursion of westerlies near the International Date Line during the middle of May associated with a Madden-Julian Oscillation (MJO) event, overall, the trades remain quite strong.




Figure 16: May 925 b zonal wind anomalies across the tropical central Pacific. Note the anomalous easterlies near the International Date Line, indicating stronger-than-normal trade winds in this area which is critical for El Niño development.

Based on this information, our best estimate is that we will likely experience neutral ENSO conditions during the 2011 hurricane season. Since we expect to continue to see a warm tropical Atlantic (discussed in detail in the next section), we believe that ENSO will not be a significant detrimental factor for this year’s hurricane season. However, there remains a need to closely monitor these conditions over the next few months.

## 7 Current Atlantic Basin Conditions

Conditions in the Atlantic remain favorable for an active season. May SSTs across the tropical Atlantic remain at above-average levels (Figure 17). The anomalously strong trades during the latter part of the winter that caused some anomalous cooling in the tropical Atlantic have now weakened (Figure 18). Sea level pressure anomalies during the month of May were also well below-average throughout the tropical Atlantic (Figure 19). Altogether, tropical Atlantic conditions currently reflect an environment conducive for an active hurricane season.




Figure 17: May 2011 SST anomaly pattern across the Atlantic Ocean.




Figure 18: May 925-mb zonal wind anomaly across the Atlantic Ocean. Note the anomalous westerly flow (e.g. weaker trades) across most of the tropical Atlantic.




Figure 19: May 2011 sea level pressure anomalies across the Atlantic. Note the anomalously low pressure throughout most of the tropical and subtropical Atlantic.

## 8 Adjusted 2011 Forecast

Table 12 shows our final adjusted early June forecast for the 2011 season which is a combination of our statistical scheme, our analog scheme and qualitative adjustments for other factors not explicitly contained in any of these schemes. Both our statistical and analog forecasts continue to call for a very active Atlantic hurricane season.

Table 12: Summary of our early June statistical forecast, our analog forecast and our adjusted final forecast for the 2011 hurricane season.

| Forecast Parameter and 1950-2000 Climatology<br>(in parentheses) | Statistical<br>Scheme | Analog<br>Scheme | Adjusted Final<br>Forecast |
|--|-----------------------|------------------|----------------------------|
| Named Storms (9.6)   | 14.1                  | 12.2             | 16                         |
| Named Storm Days (49.1)  | 77.9                  | 71.3             | 80                         |
| Hurricanes (5.9)   | 8.7                   | 7.6              | 9                          |
| Hurricane Days (24.5)  | 38.6                  | 32.4             | 35                         |
| Major Hurricanes (2.3)   | 4.5                   | 4.0              | 5                          |
| Major Hurricane Days (5.0)                                       | 11.6                  | 8.7              | 10                         |
| Accumulated Cyclone Energy Index (96.1)                          | 161                   | 139              | 160                        |
| Net Tropical Cyclone Activity (100%)                             | 172                   | 147              | 175                        |

## 9 Landfall Probabilities for 2011

A significant focus of our recent research involves efforts to develop forecasts of the probability of hurricane landfall along the U.S. coastline and in the Caribbean. Whereas individual hurricane landfall events cannot be forecast months in advance, the total seasonal probability of landfall can be forecast with statistical skill. With the observation that landfall is a function of varying climate conditions, a probability specification has been developed through statistical analyses of all U.S. hurricane and named storm landfall events during the 20<sup>th</sup> century (1900-1999). Specific landfall probabilities can be given for all tropical cyclone intensity classes for a set of distinct U.S. coastal regions.

Net landfall probability is shown linked to the overall Atlantic basin Net Tropical Cyclone activity (NTC; see Table 13). NTC is a combined measure of the year-to-year mean of six indices of hurricane activity, each expressed as a percentage difference from the long-term average. Long-term statistics show that, on average, the more active the overall Atlantic basin hurricane season is, the greater the probability of U.S. hurricane landfall.

Table 13: NTC activity in any year consists of the seasonal total of the following six parameters expressed in terms of their long-term averages. A season with 10 NS, 50 NSD, 6 H, 25 HD, 3 MH, and 5 MHD would then be the sum of the following ratios:  $10/9.6 = 104$ ,  $50/49.1 = 102$ ,  $6/5.9 = 102$ ,  $25/24.5 = 102$ ,  $3/2.3 = 130$ ,  $5/5.0 = 100$ , divided by six, yielding an NTC of 107.

| 1950-2000 Average             |      |
|-------------------------------|------|
| 1) Named Storms (NS)          | 9.6  |
| 2) Named Storm Days (NSD)     | 49.1 |
| 3) Hurricanes (H)             | 5.9  |
| 4) Hurricane Days (HD)        | 24.5 |
| 5) Major Hurricanes (MH)      | 2.3  |
| 6) Major Hurricane Days (MHD) | 5.0  |

Table 14 lists strike probabilities for the 2011 hurricane season for different TC categories for the entire U.S. coastline, the Gulf Coast and the East Coast including the Florida peninsula. We also issue probabilities for various islands and landmasses in the Caribbean and in Central America. Note that Atlantic basin NTC activity in 2011 is expected to be well above its long-term average of 100, and therefore, landfall probabilities are above their long-term average.

As an example we find that the probability of Florida being hit by a major (Cat 3-4-5) hurricane this year is 34% which is substantially higher than the yearly climatological average of 21%.

South Florida is much more prone to being impacted by a hurricane on an individual-year basis compared with northeast Florida. For instance, the probability of Miami-Dade County being impacted by hurricane-force wind gusts this year is 19%. For Duval County, the probability of being impacted by hurricane-force wind gusts is only 5%. However, considering a 50-year period, the probability of Duval County experiencing hurricane-force wind gusts is 75%.

For the island of Puerto Rico, the probability of a named storm, hurricane and major hurricane tracking within 50 miles of the island this year is 50%, 26%, and 8%, respectively.

Table 14: Estimated probability (expressed in percent) of one or more landfalling tropical storms (TS), category 1-2 hurricanes (HUR), category 3-4-5 hurricanes, total hurricanes and named storms along the entire U.S. coastline, along the Gulf Coast (Regions 1-4), and along the Florida Peninsula and the East Coast (Regions 5-11) for 2011. Probabilities of a tropical storm, hurricane and major hurricane tracking into the Caribbean are also provided. The long-term mean annual probability of one or more landfalling systems during the last 100 years is given in parentheses.

| Region                                 | TS        | Category 1-2 HUR | Category 3-4-5 HUR | All HUR   | Named Storms |
|--|-----------|------------------|--------------------|-----------|--------------|
| Entire U.S. (Regions 1-11)             | 94% (79%) | 86% (68%)        | 72% (52%)          | 96% (84%) | 99% (97%)    |
| Gulf Coast (Regions 1-4)               | 79% (59%) | 62% (42%)        | 47% (30%)          | 80% (60%) | 96% (83%)    |
| Florida plus East Coast (Regions 5-11) | 71% (50%) | 64% (44%)        | 48% (31%)          | 81% (61%) | 94% (81%)    |
| Caribbean (10-20°N, 60-88°W)           | 95% (82%) | 77% (57%)        | 61% (42%)          | 91% (75%) | 99% (96%)    |

## **10 Has Global Warming Been Responsible for the Recent Large Upswing (since 1995) in Atlantic Basin Major Hurricanes?**

### **A. BACKGROUND**

The U.S. landfall of major hurricanes Dennis, Katrina, Rita and Wilma in 2005 and the four Southeast landfalling hurricanes of 2004 – Charley, Frances, Ivan and Jeanne, raised

questions about the possible role that global warming played in those two unusually destructive seasons. In addition, three category 2 hurricanes (Dolly, Gustav and Ike) pummeled the Gulf Coast in 2008 causing considerable devastation. Some researchers have tried to link the rising CO<sub>2</sub> levels with SST increases during the late 20<sup>th</sup> century and say that this has brought on higher levels of hurricane intensity.

These speculations that hurricane intensity has increased due to CO<sub>2</sub> increases have been given much media attention; however, we believe that they are not valid, given current observational data.

There has, however, been a large increase in Atlantic basin major hurricane activity since 1995 in comparison with the prior 16-year period of 1979-1994 (Figure 20) as well as the prior quarter-century period of 1970-1994. It has been tempting for many who do not have a strong background of hurricane information to jump on this recent increase in major hurricane activity as strong evidence of a human influence on hurricanes. It should be noted, however, that the last 16-year active major hurricane period of 1995-2010 has not been more active than the earlier 16-year period of 1949-1964 when the Atlantic Ocean circulation conditions were similar to what has been observed during the last 16 years. These earlier active conditions occurred even though atmospheric CO<sub>2</sub> amounts were lower during the earlier period.






Figure 20: The tracks of major (Category 3-4-5) hurricanes during the 16-year period of 1995-2010 when the Atlantic thermohaline circulation (THC) was strong versus the prior 16-year period of 1979-1994 when the THC was weak. Note that there were approximately 2.5 times as many major hurricanes when the THC was strong as when it was weak.

Table 15 shows how large Atlantic basin hurricane variations are between strong and weak THC periods. Note especially how large the ratio is for major hurricane days (3.7) during strong vs. weak THC periods. Normalized U.S. hurricane damage studies by Pielke and Landsea (1998) and Pielke et al. (2008) show that landfalling major hurricanes account on average for about 80-85 percent of all hurricane-related destruction. This occurs even though these major hurricanes make up only 20-25 percent of named storms.

Although global surface temperatures increased during the late 20<sup>th</sup> century, there is no reliable data to indicate increased hurricane frequency or intensity in any of the globe's other tropical cyclone basins since 1972. Global Accumulated Cyclone Energy (ACE), defined as the sum of the square of a named storm's maximum wind speed (in  $10^4$  knots<sup>2</sup>) for each 6-hour period of its existence, shows significant year-to-year and decadal variability over the past forty years but no increasing trend (Figure 21). Similarly,

Klotzbach (2006) found no significant change in global TC activity during the period from 1986-2005.

Table 15: Comparison of Atlantic annual basin hurricane activity in two 16-year periods when the Atlantic Ocean THC (or AMO) was strong versus an intermediate period (1970-1994) when the THC was weak.

|                                 | THC    | SST (10-15°N;<br>70-40°W)     | Avg.<br>CO <sub>2</sub><br>ppm | NS   | NSD  | H   | HD   | MH  | MHD | ACE | NTC |
|---------------------------------|--------|-------------------------------|--------------------------------|------|------|-----|------|-----|-----|-----|-----|
| 1949-1964 (16 years)            | Strong | 27.93                         | 319                            | 10.1 | 54.1 | 6.5 | 29.9 | 3.8 | 9.5 | 121 | 133 |
| 1970-1994 (25 years)            | Weak   | 27.60                         | 345                            | 9.3  | 41.9 | 5.0 | 16.0 | 1.5 | 2.5 | 68  | 75  |
| 1995-2010 (16 years)            | Strong | 28.02                         | 373                            | 14.6 | 74.1 | 7.8 | 32.0 | 3.8 | 9.4 | 140 | 153 |
| Annual Ratio<br>Strong/Weak THC |        | $\Delta 0.35^{\circ}\text{C}$ | $\sim 0$                       | 1.3  | 1.5  | 1.4 | 1.9  | 2.5 | 3.7 | 1.9 | 1.9 |




Figure 21: Northern Hemisphere and global Accumulated Cyclone Energy (ACE) over the period from December 1971-April 2011. Figure has been adapted from Ryan Maue.

**Causes of the Upswing in Atlantic Major Hurricane Activity since 1995.** The Atlantic Ocean has a strong multi-decadal signal in its hurricane activity which is likely due to multi-decadal variations in the strength of the THC (Figure 22). The oceanic and

atmospheric response to the THC is often referred to as the Atlantic Multi-decadal Oscillation (AMO). We use the THC and AMO interchangeably throughout the remainder of this discussion. The strength of the THC can never be directly measured, but it can be diagnosed, as we have done, from the magnitude of the sea surface temperature anomaly (SSTA) in the North Atlantic (Figure 23) combined with the sea level pressure anomaly (SLPA) in the Atlantic between the latitude of the equator and 50°N (Klotzbach and Gray 2008).

The THC (or AMO) is strong when there is an above-average poleward advection of warm low-latitude waters to the high latitudes of the North Atlantic. This water can then sink to deep levels when it reaches the far North Atlantic in a process known as deep water formation. The water then moves southward at deep levels in the ocean. The amount of North Atlantic water that sinks is proportional to the water’s density which is determined by its salinity content as well as its temperature. Salty water is denser than fresh water especially at water temperatures near freezing. There is a strong association between North Atlantic SSTA and North Atlantic salinity (Figure 24). High salinity implies higher rates of North Atlantic deep water formation (or subsidence) and thus a stronger flow of upper level warm water from lower latitudes as replacement. See the papers by Gray et al. (1999), Goldenberg et al. (2001), and Grossman and Klotzbach (2009) for more discussion.




Figure 22: Illustration of strong (top) and weak (bottom) phases of the THC or AMO.




Figure 23: Long-period portrayal (1878-2006) of North Atlantic sea surface temperature anomalies (SSTA). The red (warm) periods are when the THC (or AMO) is stronger than average and the blue periods are when the THC (or AMO) is weaker than average.




Figure 24: Illustration of the strong association of yearly average North Atlantic SSTA and North Atlantic salinity content between 1958 and 2004.

## B. WHY CO<sub>2</sub> INCREASES ARE NOT RESPONSIBLE FOR ATLANTIC SST AND HURRICANE ACTIVITY INCREASES

Theoretical considerations do not support a close relationship between SSTs and hurricane intensity. In a global warming world, the atmosphere's upper air temperatures will warm or cool in unison with longer-period SST changes. Vertical lapse rates will thus not be significantly altered in a somewhat warmer or somewhat cooler tropical oceanic environment. We have no plausible physical reasons for believing that Atlantic hurricane frequency or intensity will significantly change if global or Atlantic Ocean temperatures were to rise by 1-2°C. Without corresponding changes in many other basic features, such as vertical wind shear or mid-level moisture, little or no additional TC activity should occur with SST increases.

**Confusing Time Scales of SST Influences.** A hurricane passing over a warmer body of water, such as the Gulf Stream, will often undergo some intensification. This is due to the sudden lapse rate increase which the hurricane's inner core experiences when it passes over warmer water. The warmer SSTs cause the hurricane's lower boundary layer temperature and moisture content to rise. While these low level changes are occurring, upper tropospheric conditions are often not altered significantly. These rapidly occurring lower- and upper-level temperature differences cause the inner-core hurricane lapse rates to increase and produce more intense inner-core deep cumulus convection. This typically causes a rapid increase in hurricane intensity. Such observations have led many observers to directly associate SST increases with greater hurricane potential intensity. This is valid reasoning for day-to-day hurricane intensity change associated with hurricanes moving over warmer or colder patches of SST. But such direct reasoning does not hold for conditions occurring in an overall climatologically warmer (or cooler) tropical oceanic environment where broad-scale global and tropical rainfall conditions are not expected to significantly vary. During long-period climate change, temperature and moisture conditions rise at both lower and upper levels. Lapse rates are little affected.

Any warming-induced increase in boundary layer temperature and moisture will be (to prevent significant global rainfall alteration) largely offset by a similar but weaker change through the deep troposphere up to about 10 km height. Upper-tropospheric changes are weaker than boundary layer changes, but they occur through a much deeper layer. These weaker and deeper compensating increases in upper-level temperature and moisture are necessary to balance out the larger increases in temperature and moisture which occur in the boundary layer. Global and tropical rainfall would be altered significantly only if broad-scale lapse rates were ever altered to an appreciable degree.

Thus, we cannot automatically assume that with warmer global SSTs that we will have more intense hurricanes due to lapse-rate alterations. We should not expect that the frequency and/or intensity of major hurricanes will necessarily change as a result of changes in global or individual storm basin SSTs. Historical evidence does not support hurricanes being less intense during the late 19<sup>th</sup> century and the early part of the 20<sup>th</sup> century when SSTs were slightly lower.

**CO<sub>2</sub> Influence on Hurricane Activity.** We have been performing research with the International Satellite Cloud Climatology Project (ISCCP) and the NOAA National Centers for Environmental Prediction/National Center for Atmospheric Research

(NCEP/NCAR) Reanalysis data sets. We have used this data to make an annual average of the global tropical (30°N-30°S; 0-360°) energy budget (Figure 25) for the years from 1984-2004. Note that the various surface and top of the atmosphere energy fluxes are very large. For the tropical surface, for instance, there are 637 Wm<sup>-2</sup> units of downward incoming solar and infrared (IR) energy. This downward energy flux is largely balanced by an upward surface energy flux of 615 Wm<sup>-2</sup> which is due to upward fluxes from IR energy, evaporated liquid water, and sensible heat. Similar large energy fluxes are present at the top of the atmosphere and within the troposphere.




Figure 25: Vertical cross-section of the annual tropical energy budget as determined from a combination of ISCCP and NCEP/NCAR Reanalysis data over the period from 1984-2004. Abbreviations are **IR** for longwave infrared radiation, **Alb** for albedo and **OLR** for outgoing longwave radiation. The tropics receive an excess of about 44 Wm<sup>-2</sup> radiation energy which is convected and exported as sensible heat to latitudes poleward of 30°. Estimates are about half (22 Wm<sup>-2</sup>) of this excess is transported by the atmosphere and the other half is transported by the oceans. Note, on the right, how small an OLR blockage has occurred up to now due to CO<sub>2</sub> increases (~ 1.3 Wm<sup>-2</sup>) and a continued small blockage of 3.7 Wm<sup>-2</sup> that will occur from a doubling of CO<sub>2</sub> by the end of this century.

It has been estimated that a doubling of CO<sub>2</sub> (from the pre-industrial period) without any feedback influences would result in a blockage of OLR to space of about 3.7 Wm<sup>-2</sup>. The currently-measured value of CO<sub>2</sub> in the atmosphere is 385 parts per million by volume

(ppmv). If we take the background pre-industrial value of CO<sub>2</sub> to be 285 ppmv, then by theory we should currently be having (from CO<sub>2</sub> increases alone) about  $(100/285)*3.7 = 1.3 \text{ Wm}^{-2}$  less OLR energy flux to space than was occurring in the mid-19<sup>th</sup> century.

This reduced OLR of  $1.3 \text{ Wm}^{-2}$  is very small in comparison with most of the other tropical energy budget exchanges. Slight changes in any of these other larger tropical energy budget components could easily negate or reverse this small CO<sub>2</sub>-induced OLR blockage. For instance, an upper tropospheric warming of about 1°C with no change in moisture would enhance OLR sufficiently that it would balance the reduced OLR influence from a doubling of CO<sub>2</sub>. Similarly, if there were a reduction of upper level water vapor such that the long wave radiation emission level to space were lowered by about 7 mb (~ 140 m), there would be an enhancement of OLR (with no change of temperature) sufficient to balance the suppression of OLR from a doubling of CO<sub>2</sub>. The  $1.3 \text{ Wm}^{-2}$  reduction in OLR we have experienced since the mid-19<sup>th</sup> century (about one-third of the way to a doubling of CO<sub>2</sub>) is very small compared with the overall  $399 \text{ Wm}^{-2}$  of solar energy impinging on the top of the tropical atmosphere and the mostly compensating  $356 \text{ Wm}^{-2}$  of OLR and albedo energy going back to space. This  $1.3 \text{ Wm}^{-2}$  energy gain is much too small to ever allow a determination of its possible influence on TC activity. Any such potential CO<sub>2</sub> influence on TC activity is deeply buried as turbulence within the tropical atmospheres' many other energy components. It is possible that future higher atmospheric CO<sub>2</sub> levels may cause a small influence on global TC activity. But any such potential influence would likely never be able to be detected, given that our current measurement capabilities only allow us to assess TC intensity to within about 5 mph.

### C. DISCUSSION

In a global warming or global cooling world, the atmosphere's upper air temperatures will warm or cool in unison with the SSTs. Vertical lapse rates will not be significantly altered. We have no plausible physical reasons for believing that Atlantic hurricane frequency or intensity will change significantly if global ocean temperatures were to continue to rise. For instance, in the quarter-century period from 1945-1969 when the globe was undergoing a weak cooling trend, the Atlantic basin experienced 80 major (Cat 3-4-5) hurricanes and 201 major hurricane days. By contrast, in a similar 25-year period from 1970-1994 when the globe was undergoing a general warming trend, there were only 38 Atlantic major hurricanes (48% as many) and 63 major hurricane days (31% as many) (Figure 26). Atlantic SSTs and hurricane activity do not follow global mean temperature trends.




Figure 26: Tracks of major (Category 3-4-5) hurricanes during the 25-year period of 1945-1969 when the globe was undergoing a weak cooling versus the 25-year period of 1970-1994 when the globe was undergoing a modest warming. CO<sub>2</sub> amounts in the later period were approximately 18 percent higher than in the earlier period. Major Atlantic hurricane activity was less than half as frequent during the latter period despite warmer global temperatures.

The most reliable long-period hurricane records we have are the measurements of US landfalling TCs since 1900 (Table 16). Although global mean ocean and Atlantic SSTs have increased by about 0.4°C between two 55-year periods (1901-1955 compared with 1956-2010), the frequency of US landfall numbers actually shows a slight downward trend for the later period. This downward trend is particularly noticeable for the US East Coast and Florida Peninsula where the difference in landfall of major (Category 3-4-5) hurricanes between the 45-year period of 1921-1965 (24 landfall events) and the 45-year period of 1966-2010 (7 landfall events) was especially large (Figure 27). For the entire United States coastline, 39 major hurricanes made landfall during the earlier 45-year period (1921-1965) compared with only 26 major hurricanes for the latter 45-year period (1966-2010). This occurred despite the fact that CO<sub>2</sub> averaged approximately 365 ppm during the latter period compared with 310 ppm during the earlier period.



Table 16: U.S. landfalling tropical cyclones by intensity during two 55-year periods.

| <i>YEARS</i>         | <i>Named Storms</i> | <i>Hurricanes</i> | <i>Major Hurricanes (Cat 3-4-5)</i> | <i>Global Temperature Increase</i> |
|----------------------|---------------------|-------------------|-------------------------------------|------------------------------------|
| 1901-1955 (55 years) | 210                 | 115               | 44                                  | +0.4°C                             |
| 1956-2010 (55 years) | 180                 | 87                | 34                                  |                                    |

We should not read too much into the four very active hurricane seasons of 2004, 2005, 2008 and 2010. The activity of these years was unusual but well within natural bounds of hurricane variation.

What made the 2004-2005 and 2008 seasons so destructive was not the high frequency of major hurricanes but the high percentage of hurricanes that were steered over the US coastline. The US hurricane landfall events of these years were primarily a result of the favorable upper-air steering currents present during these years.




Figure 27: Contrast of tracks of East Coast and Florida Peninsula major landfalling hurricanes during the 45-year period of 1921-1965 versus the most recent 45-year period of 1966-2010.

Although 2005 had a record number of TCs (28 named storms), this should not be taken as an indication of something beyond natural processes. There have been several other

years with comparable hurricane activity to 2005. For instance, 1933 had 21 named storms in a year when there was no satellite or aircraft data. Records of 1933 show all 21 named storms had tracks west of 60°W where surface observations were more plentiful. If we eliminate all of the named storms of 2005 whose tracks were entirely east of 60°W and therefore may have been missed given the technology available in 1933, we reduce the 2005 named storm total by seven (to 21) – the same number as was observed in 1933.

Utilizing the National Hurricane Center's best track database of hurricane records back to 1875, six previous seasons had more hurricane days than the 2005 season. These years were 1878, 1893, 1926, 1933, 1950 and 1995. Also, five prior seasons (1893, 1926, 1950, 1961 and 2004) had more major hurricane days. Although the 2005 hurricane season was certainly one of the most active on record, it was not as much of an outlier as many have indicated.

We believe that the Atlantic basin remains in an active hurricane cycle associated with a strong THC. This active cycle is expected to continue for another decade or two at which time we should enter a quieter Atlantic major hurricane period like we experienced during the quarter-century periods of 1970-1994 and 1901-1925. Atlantic hurricanes go through multi-decadal cycles. Cycles in Atlantic major hurricanes have been observationally traced back to the mid-19<sup>th</sup> century. Changes in the THC (or AMO) have been inferred from Greenland paleo ice-core temperature measurements going back thousands of years. These changes are natural and have nothing to do with human activity.

## **11 Forthcoming Updated Forecasts of 2011 Hurricane Activity**

We will be issuing a final seasonal updates of our 2011 Atlantic basin hurricane forecast on **Wednesday, 3 August**. We will also be issuing two-week forecasts for Atlantic TC activity during the climatological peak of the season from August-October. A late-season forecast for the Caribbean will be issued in early October. A verification and discussion of all 2011 forecasts will be issued in late November 2011. All of these forecasts will be available on the web at: <http://hurricane.atmos.colostate.edu/Forecasts>.

## **12 Acknowledgments**

Besides the individuals named on page 5, there have been a number of other meteorologists that have furnished us with data and given valuable assessments of the current state of global atmospheric and oceanic conditions. These include Brian McNoldy, Art Douglas, Ray Zehr, Mark DeMaria, Todd Kimberlain, Paul Roundy and Amato Evan. In addition, Barbara Brumit and Amie Hedstrom have provided excellent manuscript, graphical and data analysis and assistance over a number of years. We have profited over the years from many in-depth discussions with most of the current and past NHC hurricane forecasters. The second author would further like to acknowledge the encouragement he has received for this type of forecasting research application from Neil Frank, Robert Sheets, Robert Burpee, Jerry Jarrell, and Max Mayfield, all former directors of the National Hurricane Center (NHC) and the current director, Bill Read.

### 13 Citations and Additional Reading

- Alexander, M. A., I. Blade, M. Newman, J. R. Lanzante, N.-C. Lau, and J. D. Scott, 2002: The atmospheric bridge: The influence of ENSO teleconnections on air-sea interaction over the global oceans. *J. Climate*, 15, 2205-2231.
- Blake, E. S., 2002: Prediction of August Atlantic basin hurricane activity. Dept. of Atmos. Sci. Paper No. 719, Colo. State Univ., Ft. Collins, CO, 80 pp.
- Blake, E. S. and W. M. Gray, 2004: Prediction of August Atlantic basin hurricane activity. *Wea. Forecasting*, 19, 1044-1060.
- Chiang, J. C. H. and D. J. Vimont, 2004: Analogous Pacific and Atlantic meridional modes of tropical atmosphere-ocean variability. *J. Climate*, 17, 4143-4158.
- DeMaria, M., J. A. Knaff and B. H. Connell, 2001: A tropical cyclone genesis parameter for the tropical Atlantic. *Wea. Forecasting*, 16, 219-233.
- Elsner, J. B., G. S. Lehmiller, and T. B. Kimberlain, 1996: Objective classification of Atlantic hurricanes. *J. Climate*, 9, 2880-2889.
- Evan, A. T., J. Dunion, J. A. Foley, A. K. Heidinger, and C. S. Velden, 2006: New evidence for a relationship between Atlantic tropical cyclone activity and African dust outbreaks, *Geophys. Res. Lett*, 33, doi:10.1029/2006GL026408.
- Goldenberg, S. B., C. W. Landsea, A. M. Mestas-Nunez, and W. M. Gray, 2001: The recent increase in Atlantic hurricane activity: Causes and Implications. *Science*, 293, 474-479.
- Goldenberg, S. B. and L. J. Shapiro, 1996: Physical mechanisms for the association of El Niño and West African rainfall with Atlantic major hurricane activity. *J. Climate*, 1169-1187.
- Gray, W. M., 1984a: Atlantic seasonal hurricane frequency: Part I: El Niño and 30 mb quasi-biennial oscillation influences. *Mon. Wea. Rev.*, 112, 1649-1668.
- Gray, W. M., 1984b: Atlantic seasonal hurricane frequency: Part II: Forecasting its variability. *Mon. Wea. Rev.*, 112, 1669-1683.
- Gray, W. M., 1990: Strong association between West African rainfall and US landfall of intense hurricanes. *Science*, 249, 1251-1256.
- Gray, W. M., C. W. Landsea, P. W. Mielke, Jr., and K. J. Berry, 1992: Predicting Atlantic seasonal hurricane activity 6-11 months in advance. *Wea. Forecasting*, 7, 440-455.
- Gray, W. M., C. W. Landsea, P. W. Mielke, Jr., and K. J. Berry, 1993: Predicting Atlantic basin seasonal tropical cyclone activity by 1 August. *Wea. Forecasting*, 8, 73-86.
- Gray, W. M., C. W. Landsea, P. W. Mielke, Jr., and K. J. Berry, 1994a: Predicting Atlantic basin seasonal tropical cyclone activity by 1 June. *Wea. Forecasting*, 9, 103-115.
- Gray, W. M., J. D. Sheaffer and C. W. Landsea, 1996: Climate trends associated with multi-decadal variability of intense Atlantic hurricane activity. Chapter 2 in "Hurricanes, Climatic Change and Socioeconomic Impacts: A Current Perspective", H. F. Diaz and R. S. Pulwarty, Eds., Westview Press, 49 pp.

- Gray, W. M., 1998: Atlantic ocean influences on multi-decadal variations in El Niño frequency and intensity. Ninth Conference on Interaction of the Sea and Atmosphere, 78th AMS Annual Meeting, 11-16 January, Phoenix, AZ, 5 pp.
- Grossmann, I. and P. J. Klotzbach, 2009: A review of North Atlantic modes of natural variability and their driving mechanisms. *J. Geophys. Res.*, 114, D24107, doi:10.1029/2009JD012728.
- Henderson-Sellers, A., H. Zhang, G. Berz, K. Emanuel, W. Gray, C. Landsea, G. Holland, J. Lighthill, S.-L. Shieh, P. Webster, and K. McGuffie, 1998: Tropical cyclones and global climate change: A post-IPCC assessment. *Bull. Amer. Meteor. Soc.*, 79, 19-38.
- Klotzbach, P. J., 2002: Forecasting September Atlantic basin tropical cyclone activity at zero and one-month lead times. Dept. of Atmos. Sci. Paper No. 723, Colo. State Univ., Ft. Collins, CO, 91 pp.
- Klotzbach, P. J., 2006: Trends in global tropical cyclone activity over the past twenty years (1986-2005). *Geophys. Res. Lett.*, 33, doi:10.1029/2006GL025881.
- Klotzbach, P. J., 2007: Revised prediction of seasonal Atlantic basin tropical cyclone activity from 1 August. *Wea. and Forecasting*, 22, 937-949.
- Klotzbach, P. J. and W. M. Gray, 2003: Forecasting September Atlantic basin tropical cyclone activity. *Wea. and Forecasting*, 18, 1109-1128.
- Klotzbach, P. J. and W. M. Gray, 2004: Updated 6-11 month prediction of Atlantic basin seasonal hurricane activity. *Wea. and Forecasting*, 19, 917-934.
- Klotzbach, P. J. and W. M. Gray, 2006: Causes of the unusually destructive 2004 Atlantic basin hurricane season. *Bull. Amer. Meteor. Soc.*, 87, 1325-1333.
- Knaff, J. A., 1997: Implications of summertime sea level pressure anomalies. *J. Climate*, 10, 789-804.
- Knaff, J. A., 1998: Predicting summertime Caribbean sea level pressure. *Wea. and Forecasting*, 13, 740-752.
- Kossin, J. P., and D. J. Vimont, 2007: A more general framework for understanding Atlantic hurricane variability and trends. *Bull. Amer. Meteor. Soc.*, 88, 1767-1781.
- Landsea, C. W., 1991: West African monsoonal rainfall and intense hurricane associations. Dept. of Atmos. Sci. Paper, Colo. State Univ., Ft. Collins, CO, 272 pp.
- Landsea, C. W., 1993: A climatology of intense (or major) Atlantic hurricanes. *Mon. Wea. Rev.*, 121, 1703-1713.
- Landsea, C. W., 2007: Counting Atlantic tropical cyclones back to 1900. *EOS*, 88, 197, 202.
- Landsea, C. W. and W. M. Gray, 1992: The strong association between Western Sahel monsoon rainfall and intense Atlantic hurricanes. *J. Climate*, 5, 435-453.
- Landsea, C. W., W. M. Gray, P. W. Mielke, Jr., and K. J. Berry, 1992: Long-term variations of Western Sahelian monsoon rainfall and intense U.S. landfalling hurricanes. *J. Climate*, 5, 1528-1534.
- Landsea, C. W., W. M. Gray, K. J. Berry and P. W. Mielke, Jr., 1996: June to September rainfall in the African Sahel: A seasonal forecast for 1996. 4 pp.
- Landsea, C. W., N. Nicholls, W.M. Gray, and L.A. Avila, 1996: Downward trends in the frequency of intense Atlantic hurricanes during the past five decades. *Geo. Res. Letters*, 23, 1697-1700.

- Landsea, C. W., R. A. Pielke, Jr., A. M. Mestas-Nunez, and J. A. Knaff, 1999: Atlantic basin hurricanes: Indices of climatic changes. *Climatic Changes*, 42, 89-129.
- Landsea, C.W. et al., 2005: Atlantic hurricane database re-analysis project. Available online at [http://www.aoml.noaa.gov/hrd/data\\_sub/re\\_anal.html](http://www.aoml.noaa.gov/hrd/data_sub/re_anal.html)
- Mielke, P. W., K. J. Berry, C. W. Landsea and W. M. Gray, 1996: Artificial skill and validation in meteorological forecasting. *Wea. Forecasting*, 11, 153-169.
- Mielke, P. W., K. J. Berry, C. W. Landsea and W. M. Gray, 1997: A single sample estimate of shrinkage in meteorological forecasting. *Wea. Forecasting*, 12, 847-858.
- Pielke, Jr. R. A., and C. W. Landsea, 1998: Normalized Atlantic hurricane damage, 1925-1995. *Wea. Forecasting*, 13, 621-631.
- Pielke, Jr. R. A., and J. Gratz, C. W. Landsea, D. Collins, and R. Masulin, 2008: Normalized hurricane damage in the United States: 1900-2005. *Nat. Haz. Rev.*, 9, 29-42, doi:10.1061/(ASCE)1527-6988(2008)9:1(29).
- Rasmusson, E. M. and T. H. Carpenter, 1982: Variations in tropical sea-surface temperature and surface wind fields associated with the Southern Oscillation/El Niño. *Mon. Wea. Rev.*, 110, 354-384.
- Seseske, S. A., 2004: Forecasting summer/fall El Niño-Southern Oscillation events at 6-11 month lead times. Dept. of Atmos. Sci. Paper No. 749, Colo. State Univ., Ft. Collins, CO, 104 pp.
- Vimont, D. J., and J. P. Kossin, 2007: The Atlantic meridional mode and hurricane activity. *Geophys. Res. Lett.*, 34, L07709, doi:10.1029/2007GL029683.
- Wheeler, M. C., and H. H. Hendon, 2004: An all-season real-time multivariate MJO index: Development of an index for monitoring and prediction. *Mon. Wea. Rev.*, 132, 1917-1932.

## 14 Verification of Previous Forecasts

Table 17: Summary verification of the authors' five previous years of seasonal forecasts for Atlantic TC activity between 2006-2010.

| 2006                          | 6 Dec. 2005  | Update<br>4 April | Update<br>31 May | Update<br>3 August | Obs.  |
|-------------------------------|--------------|-------------------|------------------|--------------------|-------|
| Hurricanes                    | 9            | 9                 | 9                | 7                  | 5     |
| Named Storms                  | 17           | 17                | 17               | 15                 | 10    |
| Hurricane Days                | 45           | 45                | 45               | 35                 | 21.25 |
| Named Storm Days              | 85           | 85                | 85               | 75                 | 52.75 |
| Major Hurricanes              | 5            | 5                 | 5                | 3                  | 2     |
| Major Hurricane Days          | 13           | 13                | 13               | 8                  | 2     |
| Net Tropical Cyclone Activity | 195          | 195               | 195              | 140                | 85    |
| 2007                          | 8 Dec. 2006  | Update<br>3 April | Update<br>31 May | Update<br>3 August | Obs.  |
| Hurricanes                    | 7            | 9                 | 9                | 8                  | 6     |
| Named Storms                  | 14           | 17                | 17               | 15                 | 15    |
| Hurricane Days                | 35           | 40                | 40               | 35                 | 12.25 |
| Named Storm Days              | 70           | 85                | 85               | 75                 | 37.75 |
| Major Hurricanes              | 3            | 5                 | 5                | 4                  | 2     |
| Major Hurricane Days          | 8            | 11                | 11               | 10                 | 6     |
| Net Tropical Cyclone Activity | 140          | 185               | 185              | 160                | 99    |
| 2008                          | 7 Dec. 2007  | Update<br>9 April | Update<br>3 June | Update<br>5 August | Obs.  |
| Hurricanes                    | 7            | 8                 | 8                | 9                  | 8     |
| Named Storms                  | 13           | 15                | 15               | 17                 | 16    |
| Hurricane Days                | 30           | 40                | 40               | 45                 | 30.50 |
| Named Storm Days              | 60           | 80                | 80               | 90                 | 88.25 |
| Major Hurricanes              | 3            | 4                 | 4                | 5                  | 5     |
| Major Hurricane Days          | 6            | 9                 | 9                | 11                 | 7.50  |
| Accumulated Cyclone Energy    | 115          | 150               | 150              | 175                | 146   |
| Net Tropical Cyclone Activity | 125          | 160               | 160              | 190                | 162   |
| 2009                          | 10 Dec. 2008 | Update<br>9 April | Update<br>2 June | Update<br>4 August | Obs.  |
| Hurricanes                    | 7            | 6                 | 5                | 4                  | 3     |
| Named Storms                  | 14           | 12                | 11               | 10                 | 9     |
| Hurricane Days                | 30           | 25                | 20               | 18                 | 12    |
| Named Storm Days              | 70           | 55                | 50               | 45                 | 30    |
| Major Hurricanes              | 3            | 2                 | 2                | 2                  | 2     |
| Major Hurricane Days          | 7            | 5                 | 4                | 4                  | 3.50  |
| Accumulated Cyclone Energy    | 125          | 100               | 85               | 80                 | 53    |
| Net Tropical Cyclone Activity | 135          | 105               | 90               | 85                 | 69    |
| 2010                          | 9 Dec. 2009  | Update<br>7 April | Update<br>2 June | Update<br>4 August | Obs.  |
| Hurricanes                    | 6-8          | 8                 | 10               | 10                 | 12    |
| Named Storms                  | 11-16        | 15                | 18               | 18                 | 19    |
| Hurricane Days                | 24-39        | 35                | 40               | 40                 | 38.50 |
| Named Storm Days              | 51-75        | 75                | 90               | 90                 | 89.50 |
| Major Hurricanes              | 3-5          | 4                 | 5                | 5                  | 5     |
| Major Hurricane Days          | 6-12         | 10                | 13               | 13                 | 11    |
| Accumulated Cyclone Energy    | 100-162      | 150               | 185              | 185                | 165   |
| Net Tropical Cyclone Activity | 108-172      | 160               | 195              | 195                | 196   |